arXiv:hep-ph/0410152v4 20 Feb 2006

Neutrino spin-flavor oscillations in rapidly varying external fields

Maxim Dvornikov
Institute of Terrestrial Magnetism, Ionosphere and
Radiowave Propagation (IZMIRAN)
142190, Troitsk, Moscow region, Russia[*]

(Dated: February 7, 2008)

We study neutrino spin-flavor oscillations in presence of rapidly varying external fields. The general formalism for the description of neutrino oscillations in arbitrary rapidly varying external fields is elaborated. We obtain the effective Hamiltonian which determines the evolution of the averaged neutrino wave function. We apply the general technique to neutrino oscillations in rapidly varying magnetic fields. The special case of the constant transversal and twisting magnetic fields is studied. We evaluate the effect of neutrino spin-flavor oscillations in rapidly varying magnetic fields of the Sun. The numerical solutions of the Schrödinger equation with the Hamiltonian accounting for the constant transversal and twisting magnetic fields are presented. We compare them with our approximate analytical transition probability formula and reveal good agreement at high frequencies of the twisting magnetic field.

PACS numbers: 14.60.Pq, 26.65.+t, 96.60.Hv
Keywords: neutrino spin-flavor oscillations, magnetic fields, solar neutrino problem

I. INTRODUCTION

For the first time the idea of neutrino oscillations was theoretically predicted in Ref. [1]. Since then many experimental and theoretical studies of neutrino oscillations have been carried out. One of the most interesting problems in neutrino physics is the solar neutrino deficit. Nowadays it is experimentally established (see, e.g., Ref. [2]) that the disappearance of solar electron neutrinos can be accounted for by the LMA-MSW solution [3, 4]. The process of neutrino oscillations is likely to be the most reliable explanation of the solar neutrino problem. There are several theoretical models of solar neutrino oscillations such as mentioned above LMA-MSW solution. However other scenarios like spin-flavor precession (see Refs. [5, 6, 7, 8, 9]) are also considered.

Recently resonant neutrino spin-flavor oscillations were studied in Ref. [10] where an attempt was made to reproduce the data of the major solar neutrino experiments using the peak profiles of the solar magnetic field. Different approach to the solar neutrino problem was made in Ref. [11]. Supposing that neutrino spin-flavor precession played a subdominant role in comparison with flavor oscillations the combined action of these two mechanisms was examined. It was also possible to impose the constraints on the characteristics of a neutrino, namely to restrict its magnetic moment, and the strength of the solar magnetic field. Solar electron neutrino transitions into a non-electron antineutrino were studied in Ref. [12]. On the basis of the resent SNO data the indication of the neutrino Majorana magnetic moment was obtained in that paper. A global analysis of the solar neutrino problem solution with help of the spin-flavor precession was given in Ref. [13]. Two flavor oscillations scenario in the optimized self-consistent magneto-hydrodynamical magnetic field profile was adopted. A careful analysis of the predicted solar neutrino fluxes was performed in Ref. [14] using up to date helioseismological data.

We studied neutrino oscillations in electromagnetic fields of various configurations in our previous works. First it is necessary to mention that the Lorentz invariant formalism for the description of neutrino spin-flavor oscillations was elaborated in Ref. [15]. It allows one to study neutrino oscillations in arbitrary electromagnetic fields. For example, neutrino oscillations in an electromagnetic wave and in a longitudinal, with respect to the neutrino momentum, magnetic field were considered in Ref. [15]. In Ref. [16] we worked out the method for the investigation of the neutrino evolution equation solution near the resonance point. This technique is of great importance when the evolution equation cannot be solved analytically. In Ref. [17] we have applied this method to the studying of the parametric resonance in neutrino oscillations in amplitude modulated electromagnetic wave. The major feature of Refs. [16, 17] was the use of the perturbation theory in solving the neutrino evolution equation. It should be noted that the perturbation theory was also used in Ref. [18] where the parametric resonance in neutrino oscillations in matter with periodically varying density was studied. In Ref. [19] we proposed the Lorentz invariant quasi-classical approach for the description of neutrino spin oscillations in arbitrary non-derivative external fields. This method was applied in Ref. [20] to the description of neutrino spin oscillations in gravitational fields.

In this paper we study neutrino oscillations in presence of general rapidly varying fields. It should be noted that the influence of rapidly varying fields on mechanical oscillations was investigated in Ref. [21]. It is also well known that there are certain analogies between mechani-

---

[*]Electronic address: maxdvo@izmiran.ru

cal and neutrino oscillations (see, for instance, Ref. [22]). In Sec. II we start from the neutrino evolution equation with the Hamiltonian which accounts for the neutrino interaction with rapidly varying fields. Note that we do not fix the explicit form of the Hamiltonian. Then we derive the new effective Hamiltonian governing the time evolution of the averaged neutrino wave function. Therefore the obtained new Hamiltonian allows one to study the neutrino conversion in presence of arbitrary rapidly varying external fields. Our result is also beyond the perturbation theory because no assumptions about the smallness of rapidly varying fields are made. Then, in Sec. III, we apply the general technique to neutrino oscillations in rapidly varying magnetic fields. We investigate the combination of constant transversal and twisting magnetic fields. The neutrino conversion in solar magnetic fields is discussed. In Sec. IV we present the numerical solutions of the Schrödinger equation for the neutrino system interacting with constant transversal and twisting magnetic fields and compare them with the approximate analytical solutions found in the present work.

II. GENERAL FORMALISM

Let us consider the evolution of the two neutrinos $\nu = (\nu_1, \nu_2)$, which can belong to different flavors and helicity states. The evolution of the system is described by the Schrödinger type differential equation,

$$i\frac{\mathrm{d}\nu}{\mathrm{d}t} = H\nu, \tag{1}$$

where the Hamiltonian H involves external fields, e.g., electromagnetic field, interaction with matter and the characteristics of the neutrinos. Here we do not specify the explicit form of the Hamiltonian but just suppose that it is decomposed into two terms,

$$H = H_0 + \mathcal{H}, \quad \mathcal{H}(t+T) = \mathcal{H}(t). \tag{2}$$

The first term in Eq. (2) – H_0, corresponds to the neutrino interaction with constant or slowly varying external fields. In presence of only this term the solution of Eq. (1) can be easily found. The solution is known to be periodical with the typical frequency $\Omega_0 \sim 1/L_{\mathrm{eff}}$, where L_{eff} is the oscillations length. The second term in Eq. (2) – $\mathcal{H}(t)$, corresponds to rapidly varying external fields. The frequency $\omega = 2\pi/T$ should be much greater than Ω_0: $\omega \gg \Omega_0$. For example, we can suppose that the components of the Hamiltonian depend on time like $\mathcal{H}(t) \sim \cos\omega t$ or $\sin\omega t$. Note that we do not make any assumptions about the strength of varying external fields.

We will seek the solution of the Eqs. (1) and (2) in the form (see also Ref. [21]),

$$\nu(t) = \nu_0(t) + \xi(t). \tag{3}$$

In Eq. (3) the function $\xi(t)$ is the small rapidly oscillating one with zero mean value. The mean value $\bar{f}$ of a function $f(t)$ is the time averaging over the period T. The function $\nu_0(t)$ in Eq. (3) is the slowly varying one during T. Substituting Eq. (3) in Eq. (1) we obtain

$$i\dot{\nu}_0 + i\dot{\xi} = H_0\nu_0 + \mathcal{H}\nu_0 + \mathcal{H}\xi + H_0\xi. \tag{4}$$

There are two groups of the terms in Eq. (4): rapidly and slowly varying ones. In order for this expression to be an identity the terms of each group in the left-handed side of the equation should be equal to the terms of the same group in the right-handed side.

First let us discuss rapidly varying terms. The function $\dot{\xi}$ standing in the left-handed side belongs to this group. Despite the function ξ is small, its derivative can be, in principle, not small. Indeed, it is proportional to ω and thus $\dot{\xi}$ is the great quantity. There are three rapidly varying terms in the right-handed side of Eq. (4), namely $\mathcal{H}\nu_0$, $\mathcal{H}\xi$ and $H_0\xi$. However the second and the third terms in this group are the small ones because they contain the small factor ξ. Thus we can drop the terms $\mathcal{H}\xi$ and $H_0\xi$ when we discuss rapidly varying terms. Finally one receives the equation for the function ξ

$$i\frac{\mathrm{d}\xi}{\mathrm{d}t} = \mathcal{H}\nu_0. \tag{5}$$

It should be noted that the terms in Eq. (4) containing ξ can contribute to the phase of the wave function if long evolution times are considered. However, when we discuss possible applications of this approach in Sec. III, only one period of the transition probability variation is taken into account. This problem is also analyzed in Sec. IV. Equation (5) can be easily integrated and we find its solution

$$\xi(t) = -i\left(\int \mathcal{H}(t)\,\mathrm{d}t\right)\nu_0(t). \tag{6}$$

Here we take into account that variations of the function ν_0 are small. Therefore Eq. (6) presents the time dependence of the function ξ.

Now let us proceed to the description of ν_0 evolution. We can omit $i\dot{\xi}$ and $\mathcal{H}\nu_0$ in Eq. (4) because we have already proved that these terms are equal [see Eq. (5)]. Averaging residual terms in Eq. (4) over the period T and allowing for

$$\overline{\dot{\nu}_0} = \dot{\nu}_0, \quad \overline{H_0\nu_0} = H_0\nu_0, \quad \overline{H_0\xi} = H_0\bar{\xi} = 0,$$

we obtain the equation for the function ν_0

$$i\frac{\mathrm{d}\nu_0}{\mathrm{d}t} = H_0\nu_0 + \overline{\mathcal{H}\xi}. \tag{7}$$

We cannot neglect the term $\overline{\mathcal{H}\xi}$ in Eq. (7) since it is the product of two periodically varying functions. The mean value of this quantity is not equal to zero. With help of Eq. (6) equation (7) can transformed into the more common form

$$i\frac{\mathrm{d}\nu_0}{\mathrm{d}t} = H_{\mathrm{eff}}\nu_0, \tag{8}$$

where

$$H_{\text{eff}} = H_0 - i\overline{\mathcal{H}\left(\int \mathcal{H}\mathrm{d}t\right)}. \tag{9}$$

In derivation of Eqs. (8) and (9) we do not make any assumptions about the smallness of the interaction described by the Hamiltonian $\mathcal{H}$. Thus our result is beyond the perturbation theory used in Refs. [16, 17, 18].

The Hamiltonian H can be always represented in the form,

$$H = (\boldsymbol{\sigma} \cdot \mathbf{h}),$$

where $\mathbf{h}$ is the arbitrary $3D$-vector. Using this representation we discuss two cases:

- $H_0 = (\boldsymbol{\sigma}_\perp \cdot \mathbf{h}_\perp)$, $\mathcal{H} = \sigma_3 h_3$,
- $\mathcal{H} = (\boldsymbol{\sigma}_\perp \cdot \mathbf{h}_\perp)$, $H_0 = \sigma_3 h_3$.

Here we introduce $2D$-vectors $\boldsymbol{\sigma}_\perp = (\sigma_1, \sigma_2)$ and $\mathbf{h}_\perp = (h_1, h_2)$.

In the former case the application of the developed in the present paper technique leads to no interesting consequences. Indeed, using Eq. (9) we find that

$$H_{\text{eff}} = (\boldsymbol{\sigma}_\perp \cdot \mathbf{h}_\perp) - i\widehat{\mathbb{1}}\overline{h_3\left(\int h_3 \mathrm{d}t\right)},$$

where $\widehat{\mathbb{1}}$ is the unit matrix. It is well known that terms in the Hamiltonian proportional to the unit matrix can be omitted. For example, if we study neutrino oscillations in matter with rapidly varying density $n(t)$, it will cause no essential effect on the oscillations process. However, if one accounts for all components of the vector $\mathbf{h}$ in $\mathcal{H}$, i.e. $\mathcal{H} = (\boldsymbol{\sigma} \cdot \mathbf{h})$, it could result in some interesting effects. This case requires special careful examination.

Now we consider the latter case. First we introduce two auxiliary $2D$-vectors $\mathbf{a} = \mathbf{h}_\perp$ and $\mathbf{b} = \int \mathbf{a}\, \mathrm{d}t$. Again using Eq. (9) we obtain that

$$\begin{aligned} H_{\text{eff}} = & \sigma_3 h_3 - i\widehat{\mathbb{1}}\overline{(\mathbf{a}\cdot\mathbf{b})} + \sigma_3\overline{(a_1 b_2 - a_2 b_1)} \\ & \to \sigma_3\left[h_3 + \overline{(a_1 b_2 - a_2 b_1)}\right]. \end{aligned}$$

Therefore the off-diagonal rapidly varying terms can shift the resonance point. It was shown in Ref. [19] that both axial-vector and tensor and pseudotensor interactions can cause the neutrino spin precession. Thus rapidly varying external fields of the mentioned above types, e.g., electromagnetic fields, interaction with moving and polarized matter etc., will nontrivially affect the neutrino oscillation process. We study below one the possible examples.

III. NEUTRINO SPIN-FLAVOR OSCILLATIONS IN RAPIDLY VARYING MAGNETIC FIELDS

In this section let us discuss the neutrino evolution in matter under the influence of a combination of the two types of magnetic fields

- Constant transversal magnetic field $\mathbf{B}_0$,
- Twisting magnetic field $\mathbf{B}(\mathbf{r})$.

Note that if a magnetic field is constant in time we may consider only its transversal component with respect to the neutrino velocity. These magnetic field configurations were studied separately in the majority of works devoted to neutrino spin-flavor oscillations. For instance, only constant transversal component of a magnetic field was taken into account in Refs. [5, 6], and the effect of only twisting magnetic field on neutrino oscillations was considered in Refs. [7, 8, 9].

In our case the Hamiltonians H_0 and $\mathcal{H}$ have the form

$$H_0 = \begin{pmatrix} V/2 & \mu B_0 \\ \mu B_0 & -V/2 \end{pmatrix}, \tag{10}$$

and

$$\mathcal{H} = \begin{pmatrix} 0 & \mu B e^{-i\omega t} \\ \mu B e^{i\omega t} & 0 \end{pmatrix}. \tag{11}$$

where μ is the neutrino magnetic moment. In Eq. (10) we introduced the quantity,

$$\frac{V}{2} = \frac{\Delta m^2}{4E}\Theta - \frac{G_F}{\sqrt{2}} n_{\text{eff}},$$

where Θ is the function of the vacuum mixing angle θ_{vac} (the explicit form of Θ for various transitions of the $\nu_{iL} \leftrightarrow \nu_{jR}$ type can be found in Ref. [9]), Δm^2 is the difference of the neutrino mass squared, E is the neutrino energy, n_{eff} is the effective matter density, ω is the frequency of the transversal magnetic field variation, G_F is the Fermi constant.

Using Eqs. (9)-(11) it is possible to derive the expression for H_{eff},

$$H_{\text{eff}} = \begin{pmatrix} V/2 - (\mu B)^2/\omega & \mu B_0 \\ \mu B_0 & -V/2 + (\mu B)^2/\omega \end{pmatrix}. \tag{12}$$

It should be noted that the parameters B and B_0 must have non zero values. However it is interesting to discuss the case $B_0 \to 0$. When a neutrino interacts only with twisting magnetic field the problem of neutrino oscillations can be solved exactly (see, e.g., Ref. [7]). One can explicitly write the formula for the transition probability

$$P(t) = \left(\frac{\mu B}{\Omega'}\right)^2 \sin^2 \Omega' t,$$

where $\Omega' = \sqrt{(V-\omega)^2/4 + (\mu B)^2}$. It can be shown that in the limit of rapidly varying twisting magnetic field the transition probability approaches to zero. Indeed

$$P(t)\big|_{\omega \gg \mu B, V} \to \left(\frac{2\mu B}{\omega}\right)^2 \sin^2\left(\frac{\omega}{2}t\right)\Bigg|_{\omega \gg \mu B, V} \to 0.$$

Thus we can see that the transition probability vanishes when a neutrino interacts only with rapidly varying twisting magnetic field. This result also follows from Eq. (12).

Since B_0 and B do not depend on time in Eq. (12), we can solve Eq. (8) for the case of the effective Hamiltonian given in Eq. (12). The transition probability is expressed in the following way,

$$P(t) = A \sin^2\left(\frac{\pi t}{L}\right), \tag{13}$$

where

$$A = \frac{(\mu B_0)^2}{\left[V/2 - (\mu B)^2/\omega\right]^2 + (\mu B_0)^2}, \tag{14}$$

and

$$\frac{\pi}{L} = \Omega = \sqrt{\left[V/2 - (\mu B)^2/\omega\right]^2 + (\mu B_0)^2}. \tag{15}$$

Thus the validity of the developed method (the frequency of the twisting magnetic field ω should be much greater than the characteristic "frequency" of the system at the absence of this field) can be expressed as the constraint,

$$\omega \gg \Omega_0 = \sqrt{(V/2)^2 + (\mu B_0)^2}.$$

Now let us discuss the resonance conditions in neutrino oscillations in rapidly varying magnetic fields. From Eqs. (13) and (14) it follows that if the condition is satisfied,

$$\frac{V}{2} \simeq \frac{(\mu B)^2}{\omega}, \tag{16}$$

then $A \simeq 1$ and the transition probability can achieve great values. This phenomenon is analogous to resonance amplification of spin-flavor oscillations. However, in our case the resonance is attained not due to the zero value of the parameter V.

We again emphasize that the strength of the twisting magnetic field B can be not small in the proposed technique. The only assumption made consists in the great value of ω. Therefore we can discuss two cases:

- $B_0 \gg B$,
- $B \gg B_0$.

In the former case the value of A in Eq. (14) is great independently of B. This case corresponds to the usual spin-flavor precession in constant transversal magnetic field. However, in the latter case the value of $A_0 = A(B = 0)$ is much less than unity. Hence the transition probability at the absence of the additional twisting magnetic field is small. In this case we can choose the parameters so [see Eq. (16)] that the amplitude of the transition probability is great. It is this situation which is of interest because the essential neutrino spin-flavor conversion is attained under the influence of compact perturbations of a magnetic field against a rather weak constant background magnetic field. Such magnetic field configurations can be created experimentally or observed in some astrophysical media. Magnetic field in perturbations can have twisting structure. From Eqs. (14)-(16) one can derive the restriction imposed on the B and B_0,



FIG. 1: Neutrino flux propagation in the magnetic fields of the Sun, $\mathbf{B}_0$ and $\mathbf{B}$ are the vectors of constant tranversal and twisting magnetic fields.

$$\left(\frac{B}{B_0}\right)^2 \gg 1. \tag{17}$$

Now we consider the possible application of the developed technique to spin-flavor neutrino oscillations in magnetic fields of the Sun. The solar magnetic field is unlikely to be only either constant transversal or twisting. Therefore we can apply the method elaborated in this paper to the description of the solar neutrino conversion. The propagation of the neutrino flux in the combination of constant transversal and twisting solar magnetic fields is schematically depicted in Fig. 1. We consider one of the possible channels of neutrino oscillations, namely $\nu_{eL} \leftrightarrow \nu_{\mu R}$ conversion. First we should estimate the parameter V in Eq. (10). In our case the function Θ and the effective matter density have the form (see, e.g., Ref. [9]),

$$\Theta = \frac{1 + \cos 2\theta_{\text{vac}}}{2}, \quad n_{\text{eff}} = \left(n_e - \frac{1}{2} n_n\right).$$

Let us discuss a neutrino with the following properties: $\Delta m^2 \sim 10^{-5}\,\text{eV}^2$, $\theta_{\text{vac}} \sim \pi/4$, which are not excluded by the modern experimental results (see, for instance, Ref. [23]). We take the neutrino energy $E \sim 10\,\text{MeV}$, which corresponds to the ^{8}B solar neutrinos. Matter is supposed to consist mainly of the hydrogen, i.e. $n_n \simeq 0$, $n_e = n_p$ (as a consequence of the system electroneutrality) and has the density $d \sim 1.4\,\text{g} \cdot \text{cm}^{-3}$, which is close to the mean density of the solar matter. For these parameters we obtain that $V/2 \sim 10^{-15}\,\text{eV}$.

Now let us evaluate the strength of the magnetic fields, B_0 and B, necessary for the 10% conversion of the initial ν_{eL} beam. We suppose that neutrinos conversion occurs along the distance $D = t \simeq 9.2 \times 10^{-2} R_\odot \simeq 3.2 \times 10^{14}\,\mathrm{eV}^{-1}$, where $R_\odot$ is the solar radius. We also suppose that neutrinos have the transitional magnetic moment $\mu = 10^{-11}\mu_\mathrm{B}$, where μ_B is the Bohr magneton. Setting $P(t) = 0.1$ in Eq. (13) we obtain $B_0 \simeq 17.6\,\mathrm{kG}$. From the condition (17) we can derive the strength of the twisting magnetic field: $B \simeq 3.2 B_0 \simeq 56.3\,\mathrm{kG}$. The resonance condition (16) is satisfied if $\omega \simeq 1.0 \times 10^{-14}\,\mathrm{eV}$. The close values of the twisting magnetic field frequencies ($3 \times 10^{-15}\,\mathrm{eV}$) were considered in Ref. [8].

Background matter perturbations can cause twisting magnetic fields generation. If a neutrino interacts with solar plasma, the frequencies of these magnetic fields can be evaluated with help of the magnetic hydrodynamics (MHD) methods. We suppose that the twisting magnetic field is induced by a MHD wave with the wave vector perpendicular to $\mathbf{B}_0$. The frequencies of such MHD waves can achieve values up to $1.5 \times 10^5\,\mathrm{eV}$ for $B_0 \simeq 17.6\,\mathrm{kG}$ (see, for instance, Ref. [24]). The frequency of the twisting magnetic field, used in the present work, satisfies this condition. It should be noticed that solar plasma fluctuations were studied in Ref. [25].

IV. NUMERICAL SIMULATIONS

The analysis carried out in the previous section shows that the influence of rapidly varying magnetic fields on neutrino oscillations is very important in various astrophysical environments, for example, in $\nu_{eL} \leftrightarrow \nu_{\mu R}$ oscillations in magnetic fields of the Sun. In order to substantiate the correctness of the approach developed in this paper for the description of this problem we obtain the numerical solutions of Eq. (1) with the Hamiltonian given in Eqs. (10) and (11). The numerical solutions account for all terms in Eq. (4). Then we get the transition probabilities on the basis of the numerical solutions. We also compare these numerical transition probabilities with our approximate formula for the transition probability [Eq. (13)] which is derived in the limit of rapidly varying Hamiltonian $\mathcal{H}$. The numerical simulation is carried out with MATLAB language for technical computing (version 6.5).

The measurable quantity of neutrino oscillations is the transition probability. The probabilities of the neutrino oscillations are shown on Figs. 2-4 versus $\tau = 1.5t$. We also compare them with the approximate analytical formula (13). We remind that we take $V/2 = 10^{-15}\,\mathrm{eV}$, $B_0 = 17.6\,\mathrm{kG}$, $B = 56.3\,\mathrm{kG}$. The resonance frequency is $\omega = 1.0 \times 10^{-14}\,\mathrm{eV}$. The magenta line corresponds to the transition probability computed with help of the numerical solution of Eq. (1) for the taken parameters. The black line represents the transition probability given in Eq. (13).

It is noticed in Sec. II that the small factors $\mathcal{H}\xi$ and $H_0\xi$ can contribute to the phase of the neutrino oscillations. However if we consider only one period of the neutrino transition probability variation, then there is rather good agreement between numerical and approximate analytical solutions. Moreover it clearly follows from Figs. 2-4 that the phase shift decreases if ω becomes greater. Therefore it can be seen in Figs. 2-4 that the numerical expression for the transition probability approaches to the approximate formula given in Eq. (13) at great frequencies of the twisting magnetic field. This comparison proves the validity of the elaborated technique.

At the end of this section we note that the transition



FIG. 2: Neutrino transition probability for $\nu_{eL} \leftrightarrow \nu_{\mu R}$ oscillations, $\omega = 1.0 \times 10^{-14}\,\mathrm{eV}$. The magenta line corresponds to the transition probability found with help of the numerical solution of Eq. (1), the black line represents the transition probability given in Eq. (13).



FIG. 3: Neutrino transition probability for $\nu_{eL} \leftrightarrow \nu_{\mu R}$ oscillations, $\omega = 2.0 \times 10^{-14}\,\mathrm{eV}$. The magenta line corresponds to the transition probability found with help of the numerical solution of Eq. (1), the black line represents the transition probability given in Eq. (13).



FIG. 4: Neutrino transition probability for $\nu_{eL} \leftrightarrow \nu_{\mu R}$ oscillations, $\omega = 4.0 \times 10^{-14}\,\text{eV}$. The magenta line corresponds to the transition probability found with help of the numerical solution of Eq. (1), the black line represents the transition probability given in Eq. (13).

probability presented in Fig. 2 corresponds to the parameters that almost coincide with the resonance ones (see Sec. III). Despite of the rather low frequency ($\omega = 1.0 \times 10^{-13}\,\text{eV}$), there is a good agreement between the predicted oscillations parameters such as amplitude of the transition probability [Eq. (14)] as well as oscillations length [Eq. (15)] and those obtained from the numerical solution. The best parameters coincidence happens within the first period of transition probability variation.

V. CONCLUSION

In conclusion we note that in this paper we have examined neutrino oscillations in general rapidly varying fields. We have started with the usual Schrödinger type evolution equation. The Hamiltonian involved both slowly and rapidly varying in time terms. Instead of solving the evolution equation directly we have derived the new effective Hamiltonian which described the evolution of the averaged neutrino wave function. This new effective Hamiltonian enabled one to study neutrino oscillations in arbitrary rapidly varying external fields. It is worth mentioning that, in contrast to Refs. [16, 17], the strength of rapidly varying fields has not been limited. Therefore the elaborated method was beyond the perturbation theory since we have not carried out any expansions over the strength of the external fields. We have examined the resonance conditions in the case of general rapidly varying external fields. It has been revealed that matter with rapidly varying density caused no effect to neutrino oscillations. We have demonstrated that rapidly varying axial-vector (e.g., interaction with moving or polarized matter), tensor and pseudotensor external fields could result in nontrivial effects in neutrino oscillations. On the basis of the derived new effective Hamiltonian we have considered neutrino spin-flavor oscillations in the combination of constant transversal and twisting magnetic fields. The new effective Hamiltonian for this system has been obtained. We have analyzed various limiting cases ($B_0 \gg B$ and $B \gg B_0$). The limit $B_0 \to 0$ has also been discussed. It has been shown that described mechanism of neutrino oscillations could be important for the neutrino spin-flavor conversion in magnetic fields of the Sun. We have evaluated the strengths of magnetic fields and frequency of the twisting magnetic field necessary for the 10% neutrino conversion from ν_{eL} to $\nu_{\mu R}$. The numerical solutions of the exact (without neglecting any terms) Schrödinger equation for the two neutrino system interacting with constant transversal and twisting magnetic fields have been found. We have compared them with the approximate analytical formula for the transition probability. The very good agreement at high oscillations frequencies has been established.

Acknowledgments

This research was supported by grant of Russian Science Support Foundation. The author is indebted to Victor Semikoz and all participants of the IZMIRAN theoretical department seminar for helpful discussions.

---

[1] B. Pontecorvo, JETP **7**, 172 (1958).
[2] S. N. Ahmed et al., Phys. Rev. Lett. **92**, 181301 (2004), nucl-ex/0309004.
[3] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978).
[4] S. P. Mikheev and A. Yu. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985).
[5] C.-S. Lim and W. J. Marciano, Phys. Rev. D **37**, 1368 (1988).
[6] E. Akhmedov, Phys. Lett. B **213**, 64 (1988).
[7] A. Y. Smirnov, Phys. Lett B **260**, 161 (1991).
[8] E. K. Akhmedov, S. T. Petcov, and A. Y. Smirnov, Phys. Rev. D **48**, 2167 (1993), hep-ph/9301211.
[9] G. G. Likhachev and A. I. Studenikin, JETP **81**, 419 (1995).
[10] B. C. Chauhan and J. Pulido, J. High Energy Phys. **0406**, 008 (2004), hep-ph/0402194.
[11] E. K. Akhmedov and J. Pulido, Phys. Lett. B **553**, 7 (2003), hep-ph/0209192.
[12] S. K. Kang and C. S. Kim, Phys. Lett. B **584**, 98 (2004), hep-ph/0403059.
[13] J. Barranco, O. Miranda, T. Rashba, V. Semikoz, and J. Valle, Phys. Rev. D **66**, 093009 (2002), hep-ph/0207326.
[14] S. Couvidat, S. Turck-Chieze, and A. G. Kosovichev,

New solar seismic models and the neutrino puzzle (2002), astro-ph/0203107.

[15] A. Egorov, A. Lobanov, and A. Studenikin, Phys. Lett. B **491**, 137 (2000), hep-ph/9910476.

[16] M. S. Dvornikov and A. I. Studenikin, Phys. At. Nucl. **64**, 1624 (2001).

[17] M. S. Dvornikov and A. I. Studenikin, Phys. At. Nucl. **67**, 719 (2004).

[18] P. M. Fishbane and S. G. Gasiorowicz, Phys. Rev. D **64**, 113017 (2001), hep-ph/0012230.

[19] M. Dvornikov and A. Studenikin, J. High Energy Phys. **09**, 016 (2002), hep-ph/0202113.

[20] M. Dvornikov, A. Grigoriev, and A. Studenikin, Int. J. Mod. Phys. D **14**, 308 (2005), hep-ph/0406114.

[21] P. L. Kapitza, Usp. Fiz. Nauk **44**, 7 (1951).

[22] S. Weinberg, Int. J. Mod. Phys. A **2**, 301 (1987).

[23] S. Eidelman et al., Phys. Lett. B **592**, 1 (2004).

[24] A. F. Alexandrov, L. S. Bogdankevich, and A. A. Rukhadze, *Foundations of Plasma Electrodynamics* (Vysshaya shkola, Moscow, 1978), pp. 105–111.

[25] C. Burgess, N. Dzhalilov, T. Rashba, V. Semikoz, and J. Valle, Mon. Not. Roy. Astron. Soc. **348**, 609 (2004), astro-ph/0304462.